

References

- Barton L, Wolf B, Rowlings D et al. (2015) Sampling frequency affects estimates of annual nitrous oxide fluxes. *Scientific Reports*, **5**, 15912.
- Bogner C, Wolf B, Schlather M, Huwe B (2007) Analysing flow patterns from dye tracer experiments in a forest soil using extreme value statistics. *European Journal of Soil Science*, **59**, 103–113.
- Brosy C, Krampf K, Zeeman M et al. (2017) Simultaneous multicopter-based air sampling and sensing of meteorological variables. *Atmospheric Measurement Techniques*, **10**, 2773–2784.
- Butterbach-Bahl K, Wolf B (2017) Greenhouse gases: Warming from freezing soils. *Nature Geoscience*.
- Chen WW, Wolf B, Yao Z et al. (2010) Annual methane uptake by typical semiarid steppe in Inner Mongolia. *Journal of Geophysical Research*, **115**, 1–10.
- Chen WW, Wolf B, Bruggemann N, Butterbach-Bahl K, Zheng XH (2011a) Annual emissions of greenhouse gases from sheepfolds in Inner Mongolia. *Plant and Soil*, **340**, 291–301.
- Chen WW, Wolf B, Zheng XH et al. (2011b) Annual methane uptake by temperate semiarid steppes as regulated by stocking rates, aboveground plant biomass and topsoil air permeability. *Global Change Biology*, **17**, 2803–2816.
- Chen WW, Wolf B, Zheng XH et al. (2013a) Carbon dioxide emission from temperate semiarid steppe during the non-growing season. *Atmospheric Environment*, **64**, 141–149.
- Chen W, Zheng X, Chen Q, Wolf B, Butterbach-Bahl K, Brüggemann N, Lin S (2013b) Effects of increasing precipitation and nitrogen deposition on CH₄ and N₂O fluxes and ecosystem respiration in a degraded steppe in Inner Mongolia, China. *Geoderma*, **192**, 335–340.
- Chen W, Zheng X, Wolf B, Yao Z, Liu C, Butterbach-Bahl K (2017) THE POTENTIAL OF SOIL CARBON DIOXIDE, METHANE AND NITROUS OXIDE EXCHANGES OF DIFFERENTLY GRAZED SEMIARID STEPPES: BASED ON SOIL CORE EXPERIMENT. *Fresenius Environmental Bulletin*, **26**, 2571–2581.
- Denk TRA, Mohn J, Decock C et al. (2017) The nitrogen cycle: A review of isotope effects and isotope modeling approaches. *Soil Biology and Biochemistry*, **105**, 121–137.
- Diaz-Pines E, Molina-Herrera S, Dannenmann M et al. (2016) Nitrate leaching and soil nitrous oxide emissions diminish with time in a hybrid poplar short-rotation coppice in southern Germany. *GCB Bioenergy*, **9**, 613–626.
- Golston LM, Tao L, Brosy C et al. (2017) Lightweight mid-infrared methane sensor for unmanned aerial systems. *Applied Physics B: Lasers and Optics*, **123**, 1–9.
- Harris E, Joss A, Emmenegger L, Kipf M, Wolf B, Mohn J, Wunderlin P (2015) Isotopic evidence for nitrous oxide production pathways in a partial nitrification-anammox reactor. *Water Research*, **83**, 258–270.
- Heil J, Wolf B, Bruggemann N, Emmenegger L, Tuzson B, Vereecken H, Mohn J (2014) Site-specific ¹⁵N isotopic signatures of abiotically produced N₂O. *Geochimica et Cosmochimica Acta*, **139**, 72–82.
- Luo GJ, Bruggemann N, Wolf B, Gasche R, Grote R, Butterbach-Bahl K (2012) Decadal variability of soil CO₂, NO, N₂O, and CH₄ fluxes at the Höglwald Forest, Germany. *Biogeosciences*, **9**, 1741–1763.
- Luo GJ, Kiese R, Wolf B, Butterbach-Bahl K (2013) Effects of soil temperature and moisture on methane uptake and nitrous oxide emissions across three different ecosystem types. *Biogeosciences*, **10**, 3205–3219.
- Mohn J, Wolf B, Toyoda S et al. (2014) Inter-Laboratory assessment of nitrous oxide isotopomer

- analysis of isotopomer analysis by isotope ratio mass spectrometry and laser spectroscopy: current status and perspectives. *Rapid Communications in Mass Spectrometry*, **28**, 1995–2007.
- Pihlatie M, Christiansen JR, Aaltonen H et al. (2013) Comparison of static chambers to measure CH₄ emissions from soils. *Agricultural and Forest Meteorology*, **171–172**, 124–136.
- Schönbach P, Wolf B, Dickhöfer U et al. (2012) Grazing effects on the greenhouse gas balance of a temperate steppe ecosystem. *Nutrient Cycling in Agroecosystems*, **93**, 357–371.
- Wen Y, Chen Z, Dannenmann M et al. (2016) Disentangling gross N₂O production and consumption in soil. *Scientific Reports*, **6**, 36517.
- Wolf B, Zheng XH, Bruggemann N et al. (2010) Grazing-induced reduction of natural nitrous oxide release from continental steppe. *Nature*, **464**, 881–884.
- Wolf B, Chen WW, Bruggemann N, Zheng XH, Pumpanen J, Butterbach-Bahl K (2011) Applicability of the soil gradient method for estimating soil-atmosphere CO₂, CH₄, and N₂O fluxes for steppe soils in Inner Mongolia. *Journal of Plant Nutrition and Soil Science*, **174**, 359–372.
- Wolf B, Kiese R, Chen WW, Grote R, Zheng XH, Butterbach-Bahl K (2012) Modeling N₂O emissions from steppe in Inner Mongolia, China, with consideration of spring thaw and grazing intensity. *Plant and Soil*, **350**, 297–310.
- Wolf B, Merbold L, Decock C et al. (2015) First on-line isotopic characterization of N₂O above intensively managed grassland. *Biogeosciences*, **12**, 2517–2531.
- Wolf B, Chwala C, Fersch B et al. (2017) The scalex campaign: Scale-crossing land surface and boundary layer processes in the TERENO-prealpine observatory. *Bulletin of the American Meteorological Society*, **98**, 1217–1234.
- Wu X, Bruggemann N, Gasche R, Shen ZY, Wolf B, Butterbach-Bahl K (2010a) Environmental controls over soil-atmosphere exchange of N₂O, NO, and CO₂ in a temperate Norway spruce forest. *Global Biogeochemical Cycles*, **24**, GB2012.
- Wu X, Yao Z, Bruggemann N et al. (2010b) Effects of soil moisture and temperature on CO₂ and CH₄ soil atmosphere exchange of various land use/cover types in a semi-arid grassland in Inner Mongolia, China. *Soil Biology & Biochemistry*, **42**, 773–787.
- Wu HH, Dannenmann M, Fanselow N et al. (2011) Feedback of grazing on gross rates of N mineralization and inorganic N partitioning in steppe soils of Inner Mongolia. *Plant and Soil*, **340**, 127–139.
- Wu HH, Dannenmann M, Wolf B, Han XG, Zheng XH, Butterbach-Bahl K (2012) Seasonality of soil microbial nitrogen turnover in continental steppe soils of Inner Mongolia. *Ecosphere*, **3**, 34.
- Yao Z, Wolf B, Chen WW et al. (2010a) Spatial variability of N₂O, CH₄ and CO₂ fluxes within the Xilin River catchment of Inner Mongolia, China: a soil core study. *Plant and Soil*, **331**, 341–359.
- Yao Z, Wu X, Wolf B et al. (2010b) Soil-atmosphere exchange potential of NO and N₂O in different land use types of Inner Mongolia as affected by soil temperature, soil moisture, freeze-thaw, and drying-wetting events. *Journal of Geophysical Research-Atmospheres*, **115**.